CERN-TH/97-110
hep-ph/9705403

arXiv:hep-ph/9705403v1 23 May 1997

Squark and Gluino Production at Hadron Colliders*

Michael Spira
CERN Theory Division, CH-1211 Geneva 23, Switzerland

Abstract

The determination of the full SUSY QCD corrections to the production of squarks and gluinos at hadron colliders is reviewed. The inclusion of the NLO corrections stabilizes the theoretical predictions of the various production cross sections significantly and leads to sizeable enhancements of the most relevant cross sections. We discuss the phenomenological consequences of the results on present and future experimental analyses. Finally we investigate the impact of the corrections on the transverse momentum and rapidity distributions.

CERN-TH/97-110
hep-ph/9705403
May 1997

---

*Invited talk given at XXXIInd Rencontre de Moriond: QCD and High Energy Hadronic Interactions, Les Arcs, France, March 22–29, 1997.

1 Introduction

The search for Higgs bosons and supersymmetric particles is among the most important endeavors of present and future high energy physics. The novel colored particles, squarks and gluinos, can be searched for at the Tevatron, a $p\bar{p}$ collider with a c.m. energy of 1.8 TeV, and the future LHC, a pp collider with a c.m. energy of 14 TeV. Until now the search at the Tevatron has set the most stringent bounds on their masses. At the 95% CL, gluinos have to be heavier than about 175 GeV, while squarks with masses below about 175 GeV have been excluded for gluino masses below ~ 300 GeV 1). In the R-parity-conserving MSSM, supersymmetric particles can only be produced in pairs. All supersymmetric particles will decay to the lightest supersymmetric particle (LSP), which is most likely to be a neutralino, stable thanks to conserved R-parity. Thus the final signatures for the production of supersymmetric particles will mainly be jets and missing transverse energy, which is carried away by the invisible neutral LSP.

Squarks and gluinos can be produced via the processes

$$p\bar{p}/pp \to \tilde{q}\tilde{q}, \tilde{q}\bar{\tilde{q}}, \tilde{g}\tilde{g}, \tilde{q}\tilde{g} + X \,. \tag{1}$$

In the following we consider the production of all squarks and gluinos except stops. We assume the squarks to be mass degenerate, which is a reasonable approximation for all squark flavors except stops, while the light quarks (u, d, s, c, b) will be treated as massless particles. The production of stop pairs requires the inclusion of mass splitting and mixing effects and will be investigated elsewhere 2). In eq. (1) a summation over all possible squark flavors and charge conjugate final states should be implicitly understood. The calculation of the LO cross sections has been performed a long time ago 3). Since the [unphysical] scale dependence of the LO quantities amounts to about 50%, the determination of the NLO corrections is necessary in order to gain a reliable theoretical prediction, which can be used in present and future experimental analyses.

2 SUSY QCD corrections

The evaluation of the full SUSY QCD corrections splits into two pieces, the virtual corrections, generated by virtual particle exchanges, and the real corrections, which originate from gluon radiation and the corresponding crossed processes with three-particle final states 4).

2.1 Virtual corrections



Figure 1: *Typical diagrams of the virtual corrections.*

The one-loop virtual corrections are built up by gluon, gluino, quark and squark exchange contributions [see Fig. 1]. They have to be contracted with the LO matrix elements. The calculation of the one-loop contributions has been performed in dimensional regularization, leading to the extraction of ultraviolet, infrared and collinear singularities as poles in $\epsilon = (4-n)/2$. For the chiral γ_5 coupling we have used the naive scheme, which is well justified in the present analysis at the one-loop level. We have explicitly checked that after summing all virtual corrections no quadratic divergences are left over, in accordance with the general property of supersymmetric theories. The renormalization has been performed by identifying the squark and gluino masses with their pole masses, and defining the strong coupling in the $\overline{\rm MS}$ scheme including five light flavors in the corresponding β function. The massive particles, i.e. squarks, gluinos and top quarks, have been decoupled by subtracting their contribution at vanishing momentum transfer 5). In dimensional regularization, there is a mismatch between the gluonic degrees of freedom [d.o.f. $= n-2$] and those of the gluino [d.o.f. $= 2$], so that SUSY is explicitly broken. In order to restore SUSY in the physical observables in the massless limit, an additional finite counter-term is required for the renormalization of the novel $\tilde{q}\tilde{g}\bar{q}$ vertex 6).

2.2 Real corrections



Figure 2: *Typical diagrams of the real corrections.*

The real corrections originate from the radiation of a gluon in all possible ways and from the crossed processes by interchanging the gluon of the final state against a light quark in the initial state. The phase-space integration of the final-state particles has been performed in $n = 4 - 2\epsilon$ dimensions, leading to the extraction of infrared and collinear singularities as poles in ϵ. In order to isolate the singularities we have introduced a cutoff Δ in the invariant mass of, say, the $\bar{\tilde{q}}g$ pair, which separates soft and hard gluon radiation. After evaluating all angular integrals and adding the virtual and real corrections, the infrared singularities cancel. The left-over collinear singularities are universal and are absorbed in the renormalization of the parton densities at NLO. We defined the parton densities in the conventional $\overline{\rm MS}$ scheme including five light flavors, i.e. the squark, gluino and top quark contributions are not included in the mass factorization. Finally we end up with an ultraviolet, infrared and collinear finite partonic cross section, which is independent of the cutoff for $\Delta \to 0$.

However, there is an additional class of physical singularities, which have to be regularized 4). In the second diagram of Fig. 2 an intermediate $\tilde{q}\tilde{g}^*$ state is produced, before the [off-shell] gluino splits into a $q\bar{\tilde{q}}$ pair. If the gluino mass is larger than the common squark mass, and the partonic c.m. energy is larger than the sum of the squark and gluino masses, the intermediate gluino can be produced on its mass-shell. Thus the real corrections to $\tilde{q}\bar{\tilde{q}}$ production contain a contribution of $\tilde{q}\tilde{g}$ production. The residue of this part corresponds to $\tilde{q}\tilde{g}$ production with the subsequent gluino decay $\tilde{g} \to \bar{\tilde{q}}q$, which is already contained in the LO cross section of $\tilde{q}\tilde{g}$ pair production, including all final-state cascade decays. Thus this term has to be subtracted in order to derive a well-defined production cross section. Analogous subtractions emerge in all reactions: if the gluino mass is larger than the squark mass, the contributions from $\tilde{g} \to \tilde{q}\bar{q}, \bar{\tilde{q}}q$ have to be subtracted, and in the reverse case the contributions of squark decays into gluinos have to subtracted.

3 Results

The hadronic cross sections can be obtained from the partonic ones by convolution with the corresponding parton densities. We have performed the numerical analysis for the Tevatron and the LHC. For the natural renormalization/factorization scale choice $Q = m$, where m denotes the average mass of the final-state SUSY particles, the SUSY QCD corrections are large and positive, increasing the total cross sections by 10–90% 4). This is shown in Fig. 3, where the K factors, defined as the ratios of the NLO and LO cross sections, are presented as a function of the corresponding SUSY particle mass for the Tevatron.

We have investigated the residual scale dependence in LO and NLO, which is presented in Fig. 4. The inclusion of the NLO corrections reduces the LO scale dependence by a factor 3–4 and reaches a typical level of $\sim 15\%$, which serves as an estimate of the remaining theoretical uncertainty 4). Moreover, the dependence on different sets of parton densities is rather weak and leads to an additional uncertainty of $\sim 10\%$ 4). In order to quantify the effect of the NLO corrections on the search for squarks and gluinos at hadron colliders, we have extracted the SUSY particle masses corresponding to several fixed values of the production cross sections. These masses are increased by 10–30 GeV at the Tevatron and by 10–50 GeV at the LHC, thus enhancing the present bounds on the squark and gluino masses significantly 4).

Finally we have evaluated the QCD-corrected transverse-momentum and rapidity distributions for all different processes. As can be inferred from Fig. 5, the modification of the normalized distributions in NLO compared to LO is less than 10% for the transverse-momentum distributions and negligible for the rapidity distributions. Thus it is a sufficient approximation to rescale the LO distributions uniformly by the K factors of the total cross sections 4).

Acknowledgements
I would like to thank W. Beenakker, R. Höpker and P.M. Zerwas for their pleasant collaboration in the presented work.

References

[1] F. Abe *et al.*, CDF-Coll., Phys. Rev. Lett. **75** (1995) 613; S. Abachi *et al.*, D0-Coll., Phys. Rev. Lett. **75** (1995) 618.

[2] W. Beenakker, M. Krämer, T. Plehn, M. Spira and P.M. Zerwas, in preparation.



Figure 3: *K factors of the different squark and gluino production cross sections at the Tevatron. Parton density: GRV(94) with $Q = m$. Top mass: $m_t = 175$ GeV.*



Figure 4: *Scale and parton density dependence of the total* $\tilde{g}\tilde{g}$ *and* $\tilde{g}\tilde{q}$ *production cross sections at the Tevatron in LO and NLO. Parton densities: GRV(94) (solid), CTEQ3 (dashed) and MRS(A') (dotted); mass parameters:* $m_{\tilde{q}} = 280$ *GeV,* $m_{\tilde{g}} = 200$ *GeV and* $m_t = 175$ *GeV.*



Figure 5: *Normalized transverse-momentum and rapidity distributions of* $p\bar{p} \to \tilde{q}\bar{\tilde{q}} + X$ *at the Tevatron in LO (dotted) and NLO (solid). Parton densities: GRV('94) with* $Q = m$*; mass parameters:* $m_{\tilde{q}} = 280$ *GeV,* $m_{\tilde{g}} = 200$ *GeV and* $m_t = 175$ *GeV.*

[3] G.L. Kane and J.P. Leveillé, Phys. Lett. **B112** (1982) 227; P.R. Harrison and C.H. Llewellyn Smith, Nucl. Phys. **B213** (1983) 223 [Err. Nucl. Phys. **B223** (1983) 542]; E. Reya and D.P. Roy, Phys. Rev. **D32** (1985) 645; S. Dawson, E. Eichten and C. Quigg, Phys. Rev. **D31** (1985) 1581; H. Baer and X. Tata, Phys. Lett. **B160** (1985) 159.

[4] W. Beenakker, R. Höpker, M. Spira and P.M. Zerwas, Phys. Rev. Lett. **74** (1995) 2905, Z. Phys. **C69** (1995) 163, and report CERN-TH/96-215, hep-ph/9610490 (to appear in Nucl. Phys. **B**).

[5] J. Collins, F. Wilczek and A. Zee, Phys. Rev. **D18** (1978) 242; W.J. Marciano, Phys. Rev. **D29** (1984) 580; P. Nason, S. Dawson and R.K. Ellis, Nucl. Phys. **B303** (1988) 607.

[6] S.P. Martin and M.T. Vaughn, Phys. Lett. **B318** (1993) 331.